



Spatio-temporal light shaping for parallel nano-biophotonics

Glückstad, Jesper; Palima, Darwin

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):

Glückstad, J., & Palima, D. (2011). *Spatio-temporal light shaping for parallel nano-biophotonics*. Abstract from NanoBioTech - Montreux, Montreux, Switzerland. <http://www.nanotech-montreux.com/index.html>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Spatio-temporal light shaping for parallel nano-biophotonics

J. Glückstad and D. Palima
DTU Fotonik, Dept. of Photonics Engineering,
Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark
jesper.gluckstad@fotonik.dtu.dk
www.ppo.dk

Spatial and temporal light modulation may be visualized as forming light shapes by altering a property of light, e.g. its intensity. From this perspective, focusing light tightly and generating ultrashort light pulses are seen, respectively, as varying the width and the thickness of light. It is obvious that a 1 μm -wide focused light indeed describes the dimension of light and, though less obvious, a 10 femtosecond pulse describes a 3 μm -thick light chunk. In this picture, the “femtosecond-barrier” in ultrashort pulses and the “diffraction barrier” in focused light are mere consequences of geometry, where the wavelength of light sets a lower limit in how small we can shape light’s thickness and width.

This light shaping process suggests a natural marriage of spatial and temporal shaping (i.e., spatio-temporal modulation). However, the spatial and temporal light sciences have traditionally followed separate tracks. Width-shaping, or spatial techniques, have mostly ignored light’s thickness (using continuous-wave lasers), while thickness-shaping, or temporal techniques, typically ignored the beam width. This disconnected spatial and temporal track also shows in our own research where we developed Generalized Phase Contrast (GPC) [1] for diverse spatial light-shaping applications. We can dynamically shape light to interact with microscopic matter and we have contributed to the field of optical micromanipulation [2]. Beyond microscopy, we can apply non-contact forces onto biological cells and light-sculpted microtools [3] using our BioPhotonics Workstation which is now on its way to commercialization.

Why do we need spatiotemporal modulation? Ultrashort pulses can unravel details of ultrafast phenomena while tightly focused light can map out features with finer spatial details. However, the thin light sheets in ultrashort pulses and the narrow light shapes in focused light are simply examples of general arbitrary light shaping. For example, focusing ultrashort pulses creates narrow and thin light that achieves precision micromachining of inorganic materials such as glass and organic materials such as polymers or biological matter. Focusing ultrashort pulses compresses light and packs high energy densities that can excite multiphoton processes having much finer features. Shaping light to create an array of such high-energy chunks enables parallel processing for getting higher throughput. Another step is to vary light’s pulsewidth (thickness) as it propagates to get maximum compression (and highest energy density) at a chosen target plane. This temporal focusing can selectively look at a defined cross-section within a sample with only minimal disturbance from other regions. It can also do plane-by-plane micromachining for faster laser processing compared to scanning a focused laser spot.

Our previous work on spatial light shaping, together with the interplay between spatial and temporal modulation, invariably provides a strong position to pursue application-oriented spatio-temporal approaches [4]. Combining our GPC technology with temporal focusing, we can precisely stimulate single neuronal processes, neurons or groups of neurons, despite the highly complex neuronal structures [5] (see Fig. 1). Our recent paper in *Nature Methods*, within the emerging field of *optogenetics*, opens the use of shaped light for electrode-free and contact-free switching of brain circuits, e.g. to probe underlying mechanisms in disorders like Alzheimers or Parkinsons.




Fig. 1: Combining our GPC spatial modulation technology with temporal focusing for optogenetics: (a) Optical layout; (b) Complex neuronal structures and the spatio-temporally shaped light for switching (adapted from [5])

References

1. J. Glückstad and D. Palima, "Generalized Phase Contrast: Applications in Optics and Photonics," *Springer*, 310 pages (2009).
2. J. Glückstad, "Sorting particles with light," *Nature Materials* 3, 9-10 (2004).
3. J. Glückstad, "Optical manipulation: Sculpting the object," *Nature Photonics* 5, 7-8 (2011).
4. D. Palima and J. Glückstad, "Multi-wavelength spatial light shaping using generalized phase contrast," *Optics express*, 16, 1331-1342 (2008).
5. E. Papagiakoumou, F. Anselmi, A. Begue, V. de Sars, J. Glückstad, E. Isacoff, V. Emiliani, "Scanless two-photon excitation of channelrhodopsin-2," *Nature Methods* 7, 848-854 (2010).